

R-parity violation: Hide & Seek

M. Hirsch¹ and W. Porod²AHEP Group, Institut de Física Corpuscular – C.S.I.C. València
Edificio Institutos de Paterna, Apt 22085, E-46071 Valencia, Spain6th June 2006**Abstract**

We point out that, if R-parity is broken spontaneously, the neutralino can decay to the final state majoron plus neutrino, which from the experimental point of view is indistinguishable from the standard missing momentum signal of supersymmetry. We identify the regions of parameter space where this decay mode is dominant and show that they are independent of R-parity conserving SUSY parameters. Thus, (a) only very weak limits on R-parity violating couplings can be derived from the observation of missing momentum events and (b) at future collider experiments huge statistics might be necessary to establish that R-parity indeed is broken. Parameter combinations which give calculated relic neutralino density larger than the measured dark matter density in case of conserved R-parity are valid points in this scenario and their phenomenology at the LHC deserves to be studied.

Is R-parity broken? Undoubtedly, a definite, affirmative answer to this question would have profound implications for both, particle physics and cosmology. Low-energy precision data can be used to derive limits on R-parity violating ($\mathcal{R}_p$) couplings [1], but experimental proof that R-parity indeed is broken can only come from the observation of a decaying lightest supersymmetric particle (LSP) at a future collider experiment.

In this paper we present and discuss a scenario with spontaneously broken R-parity where the model can be easily confused with the usual Minimal Supersymmetric Standard Model (MSSM) with conserved R-parity despite the fact that the LSP decays within a typical detector of present and future high energy experiments. We concentrate here on neutral LSPs as this is the case which has been studied mostly by experimental groups. However, in principle the LSP could be charged and quasi-stable on collider time scales and, thus, be observable.

Suppose supersymmetric particles exist with masses, such that they can be produced at the LHC. To obtain evidence that the LSP decays, at least the following two conditions need to be met: (i) The decay length should not be (much) larger than the size of a typical detector. And, (ii) a minimum percentage of LSP decays, the exact number depending on

¹mahirsch@ific.uv.es²porod@ific.uv.es

the SUSY spectrum and the luminosity of the collider, of course, should lead to visible final states. It is straightforward to estimate that neutralino ($\tilde{\chi}^0$) LSP decay lengths, for typical SUSY masses in the vicinity of $\mathcal{O}(100)$ GeV, drop below $L \sim 1$ m for $\mathcal{R}_p$ couplings larger than - very roughly - $\mathcal{O}(10^{-6})$. Condition (ii), however, is more subtle. Consider the lepton number violating part ³ of the *explicit* $\mathcal{R}_p$ superpotential [3]

$$\mathcal{W} = \epsilon_i \hat{L}_i \hat{H}_u + \lambda_{ijk} \hat{L}_i \hat{L}_j \hat{E}_k^c + \lambda'_{ijk} \hat{L}_i \hat{Q}_j \hat{D}_k^c \quad (1)$$

together with the corresponding bilinear soft terms,

$$V_{\text{soft}} = \epsilon_i B_i \tilde{L}_i H_u. \quad (2)$$

For trilinear $\mathcal{R}_p$, final states of $\tilde{\chi}^0$ decays always involve either two quarks and/or charged leptons. In bilinear $\mathcal{R}_p$, scalar neutrino vacuum expectations values (vevs) induce couplings such as $C_{\tilde{\chi}^0 Z^0 \nu}$ and $C_{\tilde{\nu} \nu \nu}$, which lead to the decay $\tilde{\chi}^0 \rightarrow 3\nu$. Numerically [4], however, the branching ratio of this mode is always $\text{Br}(\tilde{\chi}^0 \rightarrow 3\nu) < 0.1$, since (a) $\text{Br}(Z^0 \rightarrow \text{inv}) \simeq 0.2$ and (b) the coupling $C_{\tilde{\chi}^0 Z^0 \nu}$ necessarily is $C_{\tilde{\chi}^0 Z^0 \nu} \simeq C_{\tilde{\chi}^0 W^\pm l^\mp}$. Thus, in explicit $\mathcal{R}_p$ condition (ii) is automatically fulfilled.

Spontaneous violation of R-parity (s- $\mathcal{R}_p$) [5, 6], on the other hand, implies the existence of a Goldstone boson, usually called the majoron (J). In s- $\mathcal{R}_p$ the neutralino can then decay according to $\tilde{\chi}^0 \rightarrow J\nu$, i.e. completely invisible. Below we will show that this decay mode can in fact easily be the dominant one, with branching ratios very close to 100 %. From the point of view of accelerator phenomenology, models with spontaneous violation of R-parity thus can resemble the MSSM with conserved R-parity and huge statistics might be necessary before it can be established that R-parity indeed is broken.

Before turning to the details of the calculation, two more comments are in order. Firstly, experiments on solar and atmospheric neutrinos have shown that neutrinos have non-zero masses and mix [7]. Many models have been proposed, which potentially can explain these data. However, the presence of lepton number violation in $\mathcal{R}_p$ inevitably implies Majorana neutrino masses. Thus, significant *upper* bounds on most $\mathcal{R}_p$ parameters follow from oscillation experiments [8] as $\mathcal{R}_p$ naturally implies a hierarchical neutrino mass spectrum. On the other hand, even the simplest bilinear $\mathcal{R}_p$ model can explain measured neutrino masses [9], as can s- $\mathcal{R}_p$ [10], if some specific combinations of $\mathcal{R}_p$ parameters obey certain *lower* bounds, see below. In our numerical calculations we will always use $\mathcal{R}_p$ parameters fitted to neutrino physics. A short comment on how our results are affected by other choices of $\mathcal{R}_p$ parameters is given near the end of this paper.

Secondly, recent WMAP data [11] have confirmed the existence of non-baryonic dark matter and measured its contribution to the energy budget of the universe with unprecedented accuracy. Because the neutralino as a dark matter candidate is lost, most advocates of SUSY declare $\mathcal{R}_p$ undesirable. However, this does not mean that in models with $\mathcal{R}_p$, the dark matter problem could not be solved. Possible DM candidates in $\mathcal{R}_p$ are (i) light

³Spontaneous $\mathcal{R}_p$ conserves baryon number. One can also use baryon parity, or any similar symmetry, to get rid of the dangerous baryon number violating terms, see [2].

gravitinos [12, 13], (ii) the axion [14, 15] or (iii) its superpartner, the axino [16], to mention a few.

Turning to the definition of our model, we note that early versions of spontaneous R_p used the left sneutrinos to break lepton number [5]. These models are now ruled out by LEP and astrophysical data [15, 17]. The model we consider [6] therefore extends the MSSM by three additional singlet superfields, namely, $\hat{\nu}^c$, $\hat{S}$ and $\hat{\Phi}$, with lepton number assignments of $L = -1, 1, 0$ respectively. The superpotential can be written as [6]

$$\begin{aligned} \mathcal{W} = & h_U^{ij} \hat{Q}_i \hat{U}_j \hat{H}_u + h_D^{ij} \hat{Q}_i \hat{D}_j \hat{H}_d + h_E^{ij} \hat{L}_i \hat{E}_j \hat{H}_d \\ & + h_\nu^i \hat{L}_i \hat{\nu}^c \hat{H}_u - h_0 \hat{H}_d \hat{H}_u \hat{\Phi} + h \hat{\Phi} \hat{\nu}^c \hat{S} + \frac{\lambda}{3!} \hat{\Phi}^3. \end{aligned} \quad (3)$$

The first three terms are the usual MSSM Yukawa terms. The terms coupling the lepton doublets to $\hat{\nu}^c$ fix lepton number. The coupling of the field $\hat{\Phi}$ with the Higgs doublets generates an effective μ -term à la Next to Minimal Supersymmetric Standard Model [18]. The last two terms, involving only singlet fields, give mass to $\hat{\nu}^c$, $\hat{S}$ and $\hat{\Phi}$, once Φ develops a vev.

For simplicity we consider only one generation of $\hat{\nu}^c$ and $\hat{S}$. Adding more generations of $\hat{\nu}^c$ or $\hat{S}$ does not add any qualitatively new features to the model. Note also, that the superpotential, Eq. (3), does not contain any terms with dimension of mass, thus potentially offering a solution to the μ -problem of supersymmetry. The soft supersymmetry breaking terms of this model can be found in [10].

The basic guiding principle in the construction of Eq. (3) is that lepton number is conserved at the level of the superpotential. However, at low energy, i.e. after electro-weak symmetry breaking, various fields acquire vevs. Besides the usual MSSM Higgs boson vevs v_D and v_U , these are $\langle \Phi \rangle = v_P/\sqrt{2}$, $\langle \hat{\nu}^c \rangle = v_R/\sqrt{2}$, $\langle \hat{S} \rangle = v_S/\sqrt{2}$ and $\langle \tilde{\nu}_i \rangle = v_{L_i}/\sqrt{2}$. Note, that $v_R \neq 0$ generates effective bilinear terms $\epsilon_i = h_\nu^i v_R/\sqrt{2}$ and that v_R , v_S and v_{L_i} violate lepton number and R-parity spontaneously.

Once R-parity is broken, Majorana neutrino masses are generated. After some algebraic manipulations one can express the effective (3,3) neutrino mass matrix in seesaw approximation as [10]

$$(\mathbf{m}_{\nu\nu}^{\text{eff}})_{ij} = a\Lambda_i\Lambda_j + b(\epsilon_i\Lambda_j + \epsilon_j\Lambda_i) + c\epsilon_i\epsilon_j, \quad (4)$$

where $\Lambda_i = \epsilon_i v_d + \mu v_{L_i}$ and $\mu = h_0 \frac{v_P}{\sqrt{2}}$. The coefficients a , b and c can be found in [10]. Equation (4) resembles closely the structure found for the neutrino mass matrix in the explicit bilinear model at the one-loop level [9]. However, with the field content of Eq. (3) the spontaneous model produces two non-zero neutrino masses already at tree-level. Note, that the coefficients b and c go to zero (and thus one of the two neutrino masses vanishes at tree-level) in the limit when either h and h_0 vanish or, equivalently, $v_P \rightarrow \infty$ or $v_R \rightarrow \infty$.

Neutrino physics puts a number of constraints on the parameters Λ_i and ϵ_i . For the current paper the exact details are unimportant, however, the most essential constraint for the following discussion is that both, $\Lambda_i/m_{\text{SUSY}}^2$ and $|\epsilon_i/\mu|$, are small. If some of the singlet fields are light, i.e. have masses in the range of $\mathcal{O}(0.1 - \text{few})$ TeV, $|\epsilon_i/\mu|$ can be as small as $|\epsilon_i/\mu| \sim 10^{-6} - 10^{-5}$ (implying $v_{L_i}/v_D \sim 10^{-6} - 10^{-4}$). On the other extreme, independent of

the singlet spectrum, $|\epsilon_i/\mu|$ can not be larger than, say, $|\epsilon_i/\mu| \sim 10^{-3}$, due to contributions from sbottom and stau loops to the neutrino mass matrix [9].

The requirement that $v_{L_i} \ll v_D$, together with the assumption of $v_{L_i}/V < 1$, can be used to find a rather simple expression for the majoron, in the basis $A'_0 = Im(H_d^0, H_u^0, \tilde{\nu}_1, \tilde{\nu}_2, \tilde{\nu}_3, \Phi, \tilde{S}, \tilde{\nu}^c)$:

$$J \simeq \left(\frac{-v_d v_L^2}{V v^2}, \frac{v_u v_L^2}{V v^2}, \frac{v_{L1}}{V}, \frac{v_{L2}}{V}, \frac{v_{L3}}{V}, 0, \frac{v_S}{V}, -\frac{v_R}{V} \right), \quad (5)$$

where $V^2 = v_S^2 + v_R^2$. We stress that neutrino data ensures $v_{L_i}/V < 1$ even for v_R as small as $\mathcal{O}(0.1)$ GeV. Thus, the majoron in our model is mainly a gauge singlet. Constraints from the measurements of the invisible width of the Z^0 boson [17] as well as all astrophysical bounds [15] are therefore easily satisfied. Note, that the majoron has no component in the Φ direction.

The coupling between the lightest neutralino, the neutrinos and the majoron in general is a complicated function of singlet and doublet parameters. However, in the limit $v_R, v_S \rightarrow \infty$ one can derive a simple approximation formula, given by

$$C_{\tilde{\chi}_1^0 \nu_k J} \simeq \frac{\tilde{\epsilon}_k}{V} N_{14} + \frac{\tilde{v}_{Lk}}{V} (g' N_{11} - g N_{12}) + \dots \quad (6)$$

Here, $\tilde{\epsilon} = U_\nu^T \cdot \vec{\epsilon}$ and $\tilde{v}_L = U_\nu^T \cdot \vec{v}_L$, where U_ν^T is the matrix which diagonalizes the neutrino mass matrix at tree-level. The dots in Eq. (6) stand for higher order terms. Equation(6) serves to show that for constant $\tilde{\epsilon}$ and $\tilde{v}_L$, $C_{\tilde{\chi}_1^0 \nu_k J} \rightarrow 0$ as v_R goes to infinity. This is as expected, since for $v_R \rightarrow \infty$ the spontaneous model approaches the explicit bilinear model. Note, that only the presence of the field $\hat{\nu}^c$ is essential for the coupling Eq. (6). If $\tilde{S}$ is absent, replace $V \rightarrow v_R$.

We now turn to the numerical results. All calculations presented below are done with the program package SPheno [19], extended to include R-parity violation. We assume that the singlet states of the model are too heavy to be produced at future collider experiments such that only MSSM particles are observed. Note, that the smallness of the $\mathcal{R}_p$ parameters guarantees that R-parity violating decay modes of all supersymmetric particles can be neglected except for the LSP and that also SUSY particle production cross sections are completely MSSM-like. We will study only points where the lightest neutralino $\tilde{\chi}_1^0$ is the LSP.

As can be seen from Eq. (6), adjusting ϵ_i and v_{L_i} from neutrino data, see Eq. (4), one expects that the most important parameters for the decay $\tilde{\chi}^0 \rightarrow J\nu$ are $|h^\nu| = \sqrt{\sum_i (h_\nu^i)^2}$ and, equivalently, v_R . Therefore, in Fig. 1(a) we plot the sum over the branching ratios of neutralino decays containing at least one visible particle in the final state as a function of $|h^\nu|$, for fixed values of all other parameters except $v_P = 10\text{--}40$ TeV. For this plot all MSSM parameters were chosen in such a way that the sparticle spectrum as well as all production cross sections very closely resemble the values of the SPA point SPSA1a' [20]. For this point one expects about $2.5 \cdot 10^7$ neutralinos at the LHC for a total luminosity of 300 fb^{-1} . As Fig. 1(a) demonstrates, the total number of events with at least some visible energy in the final state drops in this example below 100 (10) for $|h^\nu| \leq 2.5 \cdot 10^{-3}$ ($7.5 \cdot 10^{-3}$). Note, that in any SUSY event there are two $\tilde{\chi}_1^0$ and that with such small branching ratios at most




Figure 1: (a) Sum over all at least partially visible decay modes of the lightest neutralino versus $|h_\nu| = \sqrt{\sum_i (h_\nu^i)^2}$ for mSUGRA parameter point SPS1a' [20]. (b) Same points, decay length versus v_R . This is the main result of the current paper.

one of them decays visibly. Therefore, establishing that the $\tilde{\chi}_1^0$ decays at the LHC becomes challenging.

For completeness, in Fig. 1(b) we plot $c\tau$ as a function of v_R for the same points as in Fig. 1(a). For SPS1a', the neutralino decays always with $L \leq 1$ cm. Other points have different upper limits, but $L \leq 1$ m always.

We have explicitly checked that choosing different MSSM parameters reproduces Fig. 1(a) for a number of standard points from refs. [21, 22], as well as for points with a calculated $\Omega_{\tilde{\chi}^0} \simeq 1$ (in case R_p were conserved). The invisible decay width of the neutralino does essentially not depend on MSSM parameters and thus is independent of Ω_{DM} obtained in conserved R-parity.

For this model to resemble the MSSM it is not only important that the SUSY decays are MSSM-like and that the LSP decays invisibly but also that the Higgs sector behaves as in the MSSM. This is potentially in conflict with an interesting feature of this model, namely




Figure 2: R_{Jb} defined as $\Gamma(S_{h^0} \rightarrow JJ)/\Gamma(S_{h^0} \rightarrow b\bar{b})$ as a function of v_R for the mSUGRA point SPS1a' [20] for two values of the coupling h , for a number of different values of v_P , $v_P = 10$ TeV (blue) to $v_P = 40$ TeV (red) in steps of 1 TeV.

the invisible decay of the lightest doublet Higgs boson into two Majorons [10]. In Fig. 2 we show the ratio $R_{Jb} = \Gamma(S_{h^0} \rightarrow JJ)/\Gamma(S_{h^0} \rightarrow b\bar{b})$ for the same points as shown in Fig. 1, i.e. $h = 0.009$, as well as for a different choice of this parameter, $h = 0.5$. Note that the latter choice leads to invisible $\tilde{\chi}^0$ decay branching ratios indistinguishable from Fig. 1. Obviously there is enough freedom in choosing the parameters such that the invisible decay of the Higgs varies over several orders of magnitude, from a completely negligible level up to branching ratios similar to the dominant $b\bar{b}$ mode. We mention that the observed 'dips' in R_{Jb} are due to a cancellation between various contributions in the $C_{h^0 JJ}$ coupling. Figure 2 demonstrates that not only the properties of the SUSY particles can be adjusted to be MSSM-like (with conserved R-parity) but also the Higgs sector.

A final comment: We have chosen s- $\mathcal{R}_p$ parameters fitted to neutrino masses. Smaller parameters $\vec{\epsilon}$ and $\vec{v}_L$ lead to too small neutrino masses, but are allowed phenomenologically. These would lead to larger decay lengths, but not to larger branching ratios into visible final states ⁴.

In summary, we have studied the decay properties of a neutralino LSP in a model with spontaneous violation of R-parity. The neutralino decays invisibly in this model in large parts of the parameter space, such that even after several years of data taking at LHC it might be experimentally challenging to establish that R-parity is broken. A first hint of a non-standard scenario might come from cosmology, as in most of MSSM parameter space the relic density of the neutralinos is too large to be compatible with astronomical observations. However, with sufficiently large statistics it should eventually be possible to discover the rare visible decays of the lightest neutralino, e.g. into $l^\pm q\bar{q}'$ final states, and thus prove that R-parity is violated. Needless to say, that such a search requires a dedicated effort by the experimentalists.

⁴Adding some trilinear coupling, essentially unconstrained by neutrino masses [8], like λ_{121} would of course lead to a larger number of visible events.

Acknowledgments

We thank R. Tomas for discussion on supernova and the derivation of bounds on model parameters. This work was supported by Spanish grant FPA2005-01269, by the European Commission Human Potential Program RTN network MRTN-CT-2004-503369. M.H. and W.P. are supported by MCyT Ramon y Cajal contracts.

References

- [1] For reviews on limits on (mostly trilinear) $\mathcal{R}_p$ couplings see, for example: H. K. Dreiner, in: *Perspectives on Supersymmetry*, Ed. by G.L. Kane, World Scientific [arXiv:hep-ph/9707435]; R. Barbier *et al.*, Phys. Rept. **420**, 1 (2005)
- [2] H. Dreiner in [1]; L. E. Ibanez and G. G. Ross, Nucl. Phys. B **368**, 3 (1992); H. K. Dreiner, C. Luhn and M. Thormeier, Phys. Rev. D **73**, 075007 (2006) [arXiv:hep-ph/0512163].
- [3] L. J. Hall and M. Suzuki, Nucl. Phys. **B231**, 419 (1984).
- [4] W. Porod, M. Hirsch, J. Romão and J. W. F. Valle, Phys. Rev. D **63**, 115004 (2001) [arXiv:hep-ph/0011248]; F. de Campos, O. J. P. Eboli, M. B. Magro, W. Porod, D. Restrepo and J. W. F. Valle, Phys. Rev. D **71**, 075001 (2005) [arXiv:hep-ph/0501153].
- [5] C. S. Aulakh and R. N. Mohapatra, Phys. Lett. B **119**, 136 (1982).
- [6] A. Masiero and J. W. F. Valle, Phys. Lett. B **251**, 273 (1990).
- [7] Y. Fukuda *et al.* [Super-Kamiokande Collaboration], Phys. Rev. Lett. **81**, 1562 (1998) [arXiv:hep-ex/9807003]; Q. R. Ahmad *et al.* [SNO Collaboration], Phys. Rev. Lett. **89**, 011301 (2002) [arXiv:nucl-ex/0204008]; K. Eguchi *et al.* [KamLAND Collaboration], Phys. Rev. Lett. **90**, 021802 (2003) [arXiv:hep-ex/0212021].
- [8] A complete list of neutrino mass constraints on trilinear $\mathcal{R}_p$ parameters can be found in: S. Rakshit, G. Bhattacharyya and A. Raychaudhuri, Phys. Rev. D **59**, 091701 (1999) [arXiv:hep-ph/9811500].
- [9] M. Hirsch *et al.*, Phys. Rev. **D62**, 113008 (2000), [hep-ph/0004115], Err-ibid. **D65**:119901,2002; ibid **D68**, 013009 (2003) [Erratum-ibid. D **71**, 059904 (2005)]
- [10] M. Hirsch, J. C. Romão, J. W. F. Valle and A. Villanova del Moral, Phys. Rev. **D70**, 073012 (2004), [hep-ph/0407269]; *ibid.* **D73**, 055007 (2006)[hep-ph/0512257]
- [11] D. N. Spergel *et al.*, arXiv:astro-ph/0603449.
- [12] F. Takayama and M. Yamaguchi, Phys. Lett. B **485**, 388 (2000) [arXiv:hep-ph/0005214].
- [13] M. Hirsch, W. Porod and D. Restrepo, JHEP **0503**, 062 (2005) [arXiv:hep-ph/0503059].

- [14] For reviews on axions in cosmology, see, for example: J. E. Kim, Phys. Rept. **150**, 1 (1987). G. G. Raffelt, Phys. Rept. **198** (1990) 1.
- [15] For a detailed discussion see, for example: G. G. Raffelt, “Stars as laboratories for fundamental physics: The astrophysics of neutrinos, axions, and other weakly interacting particles,” Chicago, USA: Univ. Pr. (1996)
- [16] E. J. Chun and H. B. Kim, Phys. Rev. D **60**, 095006 (1999); M. Hirsch, Proc. HEP2005, Lisboa, Portugal, 21-27 Jul 2005, [http://pos.sissa.it/PoS\(HEP2005\)343](http://pos.sissa.it/PoS(HEP2005)343)
- [17] S. Eidelman *et al.* [Particle Data Group], Phys. Lett. B **592**, 1 (2004).
- [18] R. Barbieri, S. Ferrara and C. A. Savoy, Phys. Lett. B **119**, 343 (1982).
- [19] W. Porod, Comput. Phys. Commun. **153**, 275 (2003) [arXiv:hep-ph/0301101].
- [20] J. A. Aguilar-Saavedra *et al.*, Eur. Phys. J. C **46**, 43 (2006) [arXiv:hep-ph/0511344].
- [21] M. Battaglia, A. De Roeck, J. R. Ellis, F. Gianotti, K. A. Olive and L. Pape, Eur. Phys. J. C **33**, 273 (2004) [arXiv:hep-ph/0306219].
- [22] A. De Roeck, J. R. Ellis, F. Gianotti, F. Moortgat, K. A. Olive and L. Pape, arXiv:hep-ph/0508198.